

New Developments in Mplus Version 7: Part 3

Bengt Muthén & Tihomir Asparouhov

Mplus
www.statmodel.com

Presentation at Utrecht University
August 2012

Table of Contents II

- Cross-Classified / Multiple Membership Applications

Table of Contents I

- 1 **Cross-Classified Analysis, Continued**
 - 2-Mode Path Analysis: Random Contexts in Gonzalez et al.
 - 2-Mode Path Analysis: Monte Carlo Simulation
 - Cross-Classified SEM
 - Monte Carlo Simulation of Cross-Classified SEM
 - Cross-Classified Models: Types Of Random Effects
 - Random Items, Generalizability Theory
 - Random Item 2-Parameter IRT: TIMMS Example
 - Random Item Rasch IRT Example
- 2 **Advances in Longitudinal Analysis**
 - BSEM for Aggressive-Disruptive Behavior in the Classroom
 - Cross-Classified Analysis of Longitudinal Data
 - Cross-Classified Monte Carlo Simulation
 - Cross-Classified Growth Modeling: UG Example 9.27
 - Cross-Classified Analysis of Aggressive-Disruptive Behavior in the Classroom

Cross-Classified Analysis, Continued

Advanced topics:

- 2-mode path analysis
- Cross-classified SEM
- Random item IRT

Gonzalez, de Boeck, Tuerlinckx (2008). A double-structure structural equation model for three-mode data. *Psychological Methods*, 13, 337-353.

- A population of situations that might elicit negative emotional responses
- 11 situations (e.g. blamed for someone else's failure after a sports match, a fellow student fails to return your notes the day before an exam, you hear that a friend is spreading gossip about you) viewed as randomly drawn from a population of situations
- 4 binary responses: Frustration, antagonistic action, irritation, anger
- n=679 high school students
- Level 2 cluster variables are situations and students
- 1 observation for each pair of clustering units

Research questions: Which of the relationships below are significant? Are the relationships the same on the situation level as on the subject level?





Figure 3. Graphical representation of the research questions. *a*, *b*, *c*, and *d* are effect parameters.



```

VARIABLE: NAMES = frust antag irrit anger student situation;
           CLUSTER = situation student;
           CATEGORICAL = frust antag irrit anger;

DATA: FILE = gonzalez.dat;

ANALYSIS: TYPE = CROSSCLASSIFIED;
           ESTIMATOR = BAYES;
           ITERATIONS = (10000);

MODEL: %WITHIN%
       irrit anger ON frust antag;
       irrit WITH anger;
       frust WITH antag;
       %BETWEEN student%
       irrit ON frust (1);
       anger ON frust (2);
       irrit ON antag (3);
       anger ON antag (4);
       irrit; anger; irrit WITH anger;
       frust; antag; frust WITH antag;
    
```

```

%BETWEEN situation%
irrit ON frust (1);
anger ON frust (2);
irrit ON antag (3);
anger ON antag (4);
irrit; anger; irrit WITH anger;
frust; antag; frust WITH antag;
OUTPUT: TECH8 TECH9 STDY;
PLOT: TYPE = PLOT2;
    
```

M is the number of cluster units for both between levels, β is the common slope, ψ is the within-level correlation, τ is the binary outcome threshold. Table gives bias (coverage).

Para	M=10	M=20	M=30	M=50	M=100
β_1	0.13(0.92)	0.05(0.89)	0.00(0.97)	0.01(0.92)	0.01(0.94)
$\psi_{2,11}$	0.11(1.00)	0.06(0.96)	0.01(0.98)	0.00(0.89)	0.02(0.95)
$\psi_{2,12}$	0.15(0.97)	0.06(0.92)	0.05(0.97)	0.03(0.87)	0.01(0.96)
τ_1	0.12(0.93)	0.01(0.93)	0.00(0.90)	0.03(0.86)	0.00(0.91)

Small biases for $M = 10$. Due to parameter equalities information is combined from both clustering levels. Adding unconstrained level 1 model: tetrachoric correlation matrix.

Cross-Classified SEM

- General SEM model: 2-way ANOVA. Y_{pijk} is the p -th variable for individual i in cluster j and cross cluster k

$$Y_{pijk} = Y_{1pijk} + Y_{2pj} + Y_{3pk}$$

- 3 sets of structural equations - one on each level

$$Y_{1ijk} = \nu + \Lambda_1 \eta_{ijk} + \varepsilon_{ijk}$$

$$\eta_{ijk} = \alpha + B_1 \eta_{ijk} + \Gamma_1 x_{ijk} + \xi_{ijk}$$

$$Y_{2j} = \Lambda_2 \eta_j + \varepsilon_j$$

$$\eta_j = B_2 \eta_j + \Gamma_2 x_j + \xi_j$$

$$Y_{3k} = \Lambda_3 \eta_k + \varepsilon_k$$

$$\eta_k = B_3 \eta_k + \Gamma_3 x_k + \xi_k$$

Cross - Classified SEM

- The regression coefficients on level 1 can be a random effects from each of the two clustering levels: combines cross-classified SEM and cross classified HLM
- Bayesian MCMC estimation: used as a frequentist estimator.
- Easily extends to categorical variables.
- ML estimation possible only when one of the two level of clustering has small number of units.

- 1 factor at the individual level and 1 factor at each of the clustering levels, 5 indicator variables on the individual level

$$y_{pijk} = \mu_p + \lambda_{1,p}f_{1,ijk} + \lambda_{2,p}f_{2,j} + \lambda_{3,p}f_{3,k} + \varepsilon_{2,pj} + \varepsilon_{3,pk} + \varepsilon_{1,pijk}$$

- M level 2 clusters. M level 3 clusters. 1 unit within each cluster intersection. More than 1 unit is possible. Zero units possible: sparse tables
- Monte Carlo simulation: Estimation takes less than 1 min per replication

Cross-Classified Models: Types Of Random Effects

- Type 1: Random slope.
%WITHIN%
 $s \mid y$ ON x ;
 s has variance on both crossed levels. Dependent variable can be within-level factor. Covariate x should be on the WITHIN = list.
- Type 2: Random loading.
%WITHIN%
 $s \mid f$ BY y ;
 s has variance on both crossed levels. f is a within-level factor. The dependent variable can be a within-level factor.
- Type 3: Crossed random loading.
%BETWEEN level2a%
 $s \mid f$ BY y ;
 s has variance on crossed level 2b and is defined on crossed level 2a. f is a level 2a factor, s is a level 2b factor. This is a way to use the interaction term $s \cdot f$.

Table: Absolute bias and coverage for cross-classified factor analysis model

Param	M=10	M=20	M=30	M=50	M=100
$\lambda_{1,1}$	0.07(0.92)	0.03(0.89)	0.01(0.95)	0.00(0.97)	0.00(0.91)
$\theta_{1,1}$	0.05(0.96)	0.00(0.97)	0.00(0.95)	0.00(0.99)	0.00(0.94)
$\lambda_{2,p}$	0.21(0.97)	0.11(0.94)	0.10(0.93)	0.06(0.94)	0.00(0.92)
$\theta_{2,p}$	0.24(0.99)	0.10(0.95)	0.04(0.92)	0.05(0.94)	0.02(0.96)
$\lambda_{3,p}$	0.45(0.99)	0.10(0.97)	0.03(0.99)	0.01(0.92)	0.03(0.97)
$\theta_{3,p}$	0.75(1.00)	0.25(0.98)	0.15(0.97)	0.12(0.98)	0.05(0.92)
μ_p	0.01(0.99)	0.04(0.98)	0.01(0.97)	0.05(0.99)	0.00(0.97)

Perfect coverage. Level 1 parameters estimated very well. Biases when the number of clusters is small $M = 10$. Weakly informative priors can reduce the bias for small number of clusters.

Cross-Classified Interaction Model: Random Items, Generalizability Theory

- Items are random samples from a population of items.
- The same or different items may be administered to individuals.
- Suited for computer generated items and adaptive testing.
- 2-parameter IRT model

$$P(Y_{ij} = 1) = \Phi(a_j\theta_i + b_j)$$

- $a_j \sim N(a, \sigma_a)$, $b_j \sim N(b, \sigma_b)$: random discrimination and difficulty parameters
- The ability parameter is $\theta_i \sim N(0, 1)$
- Cross-classified model. Nested within items and individuals. 1 or 0 observation in each cross-classified cell.
- Interaction of two latent variables: a_j and θ_i : Type 3 crossed random loading
- The model has only 4 parameters - much more parsimonious than regular IRT models.

Random Item 2-Parameter IRT Model Setup

VARIABLE:

NAMES = u item individual;
 CLUSTER = item individual;
 CATEGORICAL = u;

ANALYSIS:

TYPE = CROSS RANDOM;
 ESTIMATOR = BAYES;

MODEL:

```
%WITHIN%

%BETWEEN individual%
s | f BY u;
f@1 u@0;
%BETWEEN item%
u s;
```

Random Item 2-Parameter IRT: TIMMS Example

- Fox (2010) Bayesian Item Response Theory. Section 4.3.3. Dutch Six Graders Math Achievement. Trends in International Mathematics and Science Study: TIMMS 2007
- 8 test items, 478 students

Table: Random 2-parameter IRT

parameter	estimate	SE
average discrimination a	0.752	0.094
average difficulty b	0.118	0.376
variation of discrimination a	0.050	0.046
variation of difficulty b	1.030	0.760

- 8 items means that there are only 8 clusters on the %between item% level and therefore the variance estimates at that level are affected by their priors. If the number of clusters is less than 10 or 20 there is prior dependence in the variance parameters.

Random Item 2-Parameter IRT: TIMMS Example, Continued

- Using factor scores estimation we can estimate item specific parameter and SE using posterior mean and posterior standard deviation.

Table: Random 2-parameter IRT item specific parameters

item	discrimination	SE	difficulty	SE
Item 1	0.797	0.11	-1.018	0.103
Item 2	0.613	0.106	-0.468	0.074
Item 3	0.905	0.148	-1.012	0.097
Item 4	0.798	0.118	-1.312	0.106
Item 5	0.538	0.099	0.644	0.064
Item 6	0.808	0.135	0.023	0.077
Item 7	0.915	0.157	0.929	0.09
Item 8	0.689	0.105	1.381	0.108

Random Item 2-Parameter IRT: TIMMS Example, Comparison With ML

Table: Random 2-parameter IRT item specific parameters

item	Bayes random discrimination	Bayes random SE	ML fixed discrimination	ML fixed SE
Item 1	0.797	0.110	0.850	0.155
Item 2	0.613	0.106	0.579	0.102
Item 3	0.905	0.148	0.959	0.170
Item 4	0.798	0.118	0.858	0.172
Item 5	0.538	0.099	0.487	0.096
Item 6	0.808	0.135	0.749	0.119
Item 7	0.915	0.157	0.929	0.159
Item 8	0.689	0.105	0.662	0.134

- Bayes random estimates are shrunk towards the mean and have smaller standard errors: shrinkage estimate

- One can add a predictor for a person's ability. For example adding gender as a predictor yields an estimate of 0.283 (0.120), saying that males have a significantly higher math mean.
- Predictors for discrimination and difficulty random effects, for example, geometry indicator.
- More parsimonious model can yield more accurate ability estimates.

- De Boeck (2008) Random item IRT models
- 24 verbal aggression items, 316 persons

$$P(Y_{ij} = 1) = \Phi(\theta_i + b_j)$$

$$b_j \sim N(b, \sigma)$$

$$\theta_i \sim N(0, \tau)$$

Table: Random Rasch IRT - variance decomposition

parameter	person τ	item σ	error
estimates(SE)	1.89(0.19)	1.46(0.53)	2.892
variance explained	30%	23%	46%

Random Item Rasch IRT Example: Simple Model Specification

MODEL:

%WITHIN%

%BETWEEN person%

y;


%BETWEEN item%

y;

Advances in Longitudinal Analysis

- An old dilemma
- Two new solutions


Categorical Items, Wide Format, Single-Level Approach



Single-level analysis with $p \times T = 2 \times 5 = 10$ variables, $T = 5$ factors.

- ML hard and impossible as T increases (numerical integration)
- WLSMV possible but hard when $p \times T$ increases and biased unless attrition is MCAR or multiple imputation is done first
- Bayes possible
- Searching for partial measurement invariance is cumbersome

Categorical Items, Long Format, Two-Level Approach




Two-level analysis with $p = 2$ variables, 1 within-factor, 2-between factors, **assuming full measurement invariance across time**.

- ML feasible
- WLSMV feasible (2-level WLSMV)
- Bayes feasible

Measurement Invariance Across Time

- Both old approaches have problems
 - Wide, single-level approach easily gets significant non-invariance and needs many modifications
 - Long, two-level approach has to assume invariance
- New solution no. 1, suitable for small to medium number of time points
 - A new wide, single-level approach where time is a fixed mode
- New solution no. 2, suitable for medium to large number of time points
 - A new long, two-level approach where time is a random mode
 - No limit on the number of time points


New Solution No. 1: Wide Format, Single-Level Approach



Single-level analysis with $p \times T = 2 \times 5 = 10$ variables, $T = 5$ factors.

- Bayes ("BSEM") using approximate measurement invariance, still identifying factor mean and variance differences across time

- New solution no. 2, time is a random mode
- A new long, two-level approach
 - Best of both worlds: Keeping the limited number of variables of the two-level approach without having to assume invariance



Two-level analysis with $p = 2$ variables.

- Bayes twolevel random approach with random measurement parameters and random factor means and variances using Type=Crossclassified: Clusters are time and person

Aggressive-Disruptive Behavior in the Classroom

Randomized field experiment in Baltimore public schools with a classroom-based intervention aimed at reducing aggressive-disruptive behavior among elementary school students (Ialongo et al., 1999).

This analysis:

- Cohort 1
- 9 binary items at 8 time points, Grade 1 - Grade 7
- $n = 1174$

Aggressive-Disruptive Behavior in the Classroom: ML vs BSEM

- Traditional ML analysis
 - 8 dimensions of integration
 - Computing time: 25:44 with Integration = Montecarlo(5000)
 - Increasing the number of time points makes ML impossible
- BSEM analysis
 - 156 parameters
 - Computing time: 4:01
 - Increasing the number of time points has relatively less impact

BSEM Input Excerpts for Aggressive-Disruptive Behavior

```

USEVARIABLES = stub1f-tease7s;
CATEGORICAL = stub1f-tease7s;
MISSING = ALL (999);
DEFINE:      CUT stub1f-tease7s (1.5);
ANALYSIS:   ESTIMATOR = BAYES;
            PROCESSORS = 2;
MODEL:      f1f by stub1f-tease1f* (lam11-lam19);
            f1s by stub1s-tease1s* (lam21-lam29);
            f2s by stub2s-tease2s* (lam31-lam39);
            f3s by stub3s-tease3s* (lam41-lam49);
            f4s by stub4s-tease4s* (lam51-lam59);
            f5s by stub5s-tease5s* (lam61-lam69);
            f6s by stub6s-tease6s* (lam71-lam79);
            f7s by stub7s-tease7s* (lam81-lam89);
            f1f@1;
    
```

BSEM Input for Aggressive-Disruptive Behavior, Continued

```

[stub1f$1-tease1f$1] (tau11-tau19);
[stub1s$1-tease1s$1] (tau21-tau29);
[stub2s$1-tease2s$1] (tau31-tau39);
[stub3s$1-tease3s$1] (tau41-tau49);
[stub4s$1-tease4s$1] (tau51-tau59);
[stub5s$1-tease5s$1] (tau61-tau69);
[stub6s$1-tease6s$1] (tau71-tau79);
[stub7s$1-tease7s$1] (tau81-tau89);
[f1f-f7s@0];
i s q | f1f@0 f1s@0.5 f2s@1.5 f3s@2.5 f4s@3.5
f5s@4.5 f6s@5.5 f7s@6.5;
q@0;
MODEL
PRIORS:   DO(1,9) DIFF(lam1#-lam8#) ~ N(0.,01);
          DO(1,9) DIFF(tau1#-tau8#) ~ N(0.,01);
OUTPUT:   TECH1 TECH8;
    
```

Estimates for Aggressive-Disruptive Behavior

	Estimate	Posterior S.D.	One-Tailed P-Value	95% C.I.		
				Lower 2.5%	Upper 2.5%	
Means						
I	0.000	0.000	1.000	0.000	0.000	
S	0.238	0.068	0.000	0.108	0.366	*
Q	-0.022	0.011	0.023	-0.043	0.000	*
Variances						
I	9.258	2.076	0.000	6.766	14.259	*
S	0.258	0.068	0.000	0.169	0.411	*
Q	0.001	0.000	0.000	0.001	0.001	

Estimates for Aggressive-Disruptive Behavior, Continued

	Estimate	Posterior S.D.	One-Tailed P-Value	95% C.I.		
				Lower 2.5%	Upper 2.5%	
F1F BY						
STUB1F	0.428	0.048	0.000	0.338	0.522	*
BKRULE1F	0.587	0.068	0.000	0.463	0.716	*
HARMO1F	0.832	0.082	0.000	0.677	0.985	*
BKTHIN1F	0.671	0.067	0.000	0.546	0.795	*
YELL1F	0.508	0.055	0.000	0.405	0.609	*
TAKEP1F	0.717	0.072	0.000	0.570	0.839	*
FIGHT1F	0.480	0.052	0.000	0.385	0.579	*
LIES1F	0.488	0.054	0.000	0.386	0.589	*
TEASE1F	0.503	0.055	0.000	0.404	0.608	*
...						
F7S BY						
STUB7S	0.360	0.049	0.000	0.273	0.458	*
BKRULE7S	0.512	0.068	0.000	0.392	0.654	*
HARMO7S	0.555	0.074	0.000	0.425	0.716	*
BKTHIN7S	0.459	0.063	0.000	0.344	0.581	*
YELL7S	0.525	0.062	0.000	0.409	0.643	*
TAKEP7S	0.500	0.069	0.000	0.372	0.634	*
FIGHT7S	0.515	0.067	0.000	0.404	0.652	*
LIES7S	0.520	0.070	0.000	0.392	0.653	*
TEASE7S	0.495	0.064	0.000	0.378	0.626	*

Displaying Non-Invariant Items: Time Points With Significant Differences Compared to the Mean ($V = 0.01$)

Item	Loading	Threshold
stub	3	1, 2, 3, 6, 8
bkrule	-	5, 8
harmo	1, 8	2, 8
bkthin	1, 2, 3, 7, 8	2, 8
yell	2, 3, 6	-
takep	1, 2, 5	1, 2, 5
fight	1, 5	1, 4
lies	-	-
tease	-	1, 4, 8

Cross-Classified Analysis of Longitudinal Data

- Observations nested within time and subject
- A large number of time points can be handled via Bayesian analysis
- A relatively small number of subjects is needed

Intensive Longitudinal Data

- Time intensive data: More longitudinal data are collected where very frequent observations are made using new tools for data collection. Walls & Schafer (2006)
- Typically multivariate models are developed but if the number of time points is large these models will fail due to too many variables and parameters involved
- Factor analysis models will be unstable over time. Is it lack of measurement invariance or insufficient model?
- Random loading and intercept models can take care of measurement and intercept invariance. A problem becomes an advantage.
- Random loading and intercept models produce more accurate estimates for the loadings and factors by borrowing information over time
- Random loading and intercept models produce more parsimonious model

Cross-Classified Analysis: Monte Carlo Simulation Generating the Data for Ex9.27

```
TITLE:          this is an example of longitudinal modeling using a
                cross-classified data approach where observations are
                nested within the cross-classification of time and subjects

MONTECARLO:
  NAMES = y1-y3;
  NOBSERVATIONS = 7500;
  NREPS = 1;
  CSIZES = 75[100(1)];! 75 subjects, 100 time points
  NCSIZE = 1[1];
  WITHIN = (level2a) y1-y3;
  SAVE = ex9.27.dat;

ANALYSIS:
  TYPE = CROSS RANDOM;
  ESTIMATOR = BAYES;
  PROCESSORS = 2;
```

MODEL

POPULATION:

```
%WITHIN%
s1-s3 | f by y1-y3;
f@1;
y1-y3*1.2; [y1-y3@0];
%BETWEEN level2a% ! across time variation
s1-s3*0.1;
[s1-s3*1.3];
y1-y3*.5;
[y1-y3@0];
%BETWEEN level2b% ! across subjects variation
f*1; [f*.5];
s1-s3@0; [s1-s3@0];
```

```
TITLE:      this is an example of a multiple indicator
            growth model with random intercepts and
            factor loadings using cross-classified
            data
DATA:      FILE = ex9.27.dat;
VARIABLE:  NAMES = y1-y3 time subject;
            USEVARIABLES = y1-y3 timescor;
            CLUSTER = subject time;
            WITHIN = timescor (time) y1-y3;
DEFINE:    timescor = (time-1)/100;
ANALYSIS:  TYPE = CROSSCLASSIFIED RANDOM;
            ESTIMATOR = BAYES;
            PROCESSORS = 2;
            BITERATIONS = (1000);
MODEL:     %WITHIN%
            s1-s3 | f BY y1-y3;
            f@1;
            s | f ON timescor; !slope growth factor s
            y1-y3; [y1-y3@0];
            %BETWEEN time% ! time variation
            s1-s3; [s1-s3]; ! random loadings
            y1-y3; [y1-y3@0]; ! random intercepts
            s@0; [s@0];
            %BETWEEN subject% ! subject variation
            f; [f]; ! intercept growth factor f
            s1-s3@0; [s1-s3@0];
            s; [s]; ! slope growth factor s
OUTPUT:    TECH1 TECH8;
```

Cross-Classified Analysis of Aggressive-Disruptive Behavior in the Classroom

- Teacher-rated measurement instrument capturing aggressive-disruptive behavior among a sample of U.S. students in Baltimore public schools (Ialongo et al., 1999).
- The instrument consists of 9 items scored as 0 (almost never) through 6 (almost always)
- A total of 1174 students are observed in 41 classrooms from Fall of Grade 1 through Grade 6 for a total of 8 time points
- The multilevel (classroom) nature of the data is ignored in the current analyses
- The item distribution is very skewed with a high percentage in the Almost Never category. The items are therefore dichotomized into Almost Never versus the other categories combined
- We analyze the data on the original scale as continuous variables and also the dichotomized scale as categorical

Aggressive-Disruptive Behavior Example Continued

- For each student a 1-factor analysis model is estimated with the 9 items at each time point
- Let Y_{pit} be the p -th item for individual i at time t
- We use cross-classified SEM. Observations are nested within individual and time.
- Although this example uses only 8 time points the models can be used with any number of time points.

Aggressive-Disruptive Behavior Example Cont'd: Model 1

- Model 1: Two-level factor model with intercept non-invariance across time

$$Y_{pit} = \mu_p + \zeta_{pt} + \xi_{pi} + \lambda_p \eta_{it} + \varepsilon_{pit}$$

- μ_p, λ_p are model parameters, $\varepsilon_{pit} \sim N(0, \theta_{w,p})$ is the residual
- $\zeta_{pt} \sim N(0, \sigma_p)$ is a random effect to accommodate intercept non-invariance across time
- To correlate the factors η_{it} within individual i

$$\eta_{it} = \eta_{b,i} + \eta_{w,it}$$

- $\eta_{b,i} \sim N(0, \psi)$ and $\eta_{w,it} \sim N(0, 1)$. The variance is fixed to 1 to identify the scale in the model
- $\xi_{pi} \sim N(0, \theta_{b,p})$ is a between level residual in the between level factor model
- Without the random effect ζ_{pt} this is just a standard two-level factor model

Aggressive-Disruptive Behavior Example Continued: Model 1 Setup

MODEL:

```
%WITHIN%  
f BY y1-y9*1 (11-19);  
f@1;  
  
%BETWEEN t1%  
y1-y9;  
  
%BETWEEN id%  
y1-y9;  
f BY y1-y9*1 (11-19);
```

Aggressive-Disruptive Behavior Example Cont'd: Model 2

- Model 2: Adding latent growth model for the factor

$$\eta_{it} = \alpha_i + \beta_i \cdot t + \eta_{w,it}$$

- $\alpha_i \sim N(0, v_\alpha)$ is the intercept and $\beta_i \sim N(\beta, v_\beta)$ is the slope. For identification purposes again $\eta_{w,it} \sim N(0, 1)$
- The model looks for developmental trajectory across time for the aggressive-disruptive behavior factor

Aggressive-Disruptive Behavior Example Continued: Model 2 Setup

MODEL:

```
! s = beta, fb = alpha  
%WITHIN%  
f BY y1-y9*1 (11-19);  
f@1;  
s | f ON time;  
  
%BETWEEN t1%  
y1-y9;  
s@0; [s@0];  
  
%BETWEEN id%  
y1-y9;  
f BY y1-y9*1 (11-19);  
s*1; [s*0];
```

Aggressive-Disruptive Behavior Example Cont'd: Model 3

- Model 3: Adding measurement non-invariance
- Replace the fixed loadings λ_p with random loadings $\lambda_{pt} \sim N(\lambda_p, w_p)$
- The random loadings accommodate measurement non-invariance across time
- All models can be estimated for continuous and categorical scale data

Aggressive-Disruptive Behavior Example Continued: Model 3 Setup

```
MODEL:
    %WITHIN%
    s1-s9 | f BY y1-y9;
    f@1;
    s | f ON time;
    %BETWEEN t1%
    y1-y9;
    f@0; [f@0];
    s@0; [s@0];
    s1-s9*1; [s1-s9*1];
    %BETWEEN id%
    y1-y9;
    f*1; [f@0];
    s*1; [s*0];
    s1-s9@0; [s1-s9@0];
```

Aggressive-Disruptive Behavior Example Continued: Model 3 Results For Continuous Analysis

Within Level					
Residual Variances					
Y1	1.073	0.022	0.000	1.029	1.119
Y9	0.630	0.014	0.000	0.604	0.658
F	1.000	0.000	0.000	1.000	1.000
Between ID Level					
Variances					
Y1	0.146	0.016	0.000	0.118	0.180
Y9	0.052	0.009	0.000	0.035	0.068
F	1.316	0.080	0.000	1.172	1.486
S	0.026	0.003	0.000	0.020	0.032
Between T1 Level					
Means					
Y1	1.632	0.120	0.000	1.377	1.885
Y9	1.232	0.096	0.000	1.044	1.420
S1	0.679	0.023	0.000	0.640	0.732
S9	0.705	0.043	0.000	0.628	0.797
Variances					
Y1	0.080	0.138	0.000	0.025	0.372
Y9	0.047	0.109	0.000	0.017	0.266
S1	0.002	0.004	0.000	0.000	0.013
S9	0.010	0.079	0.000	0.003	0.052

Aggressive-Disruptive Behavior Example Cont'd: Model 4

- Model 4: Adding measurement non-invariance also across individuals
- Replace the loadings λ_{pt} with random loadings

$$\lambda_{pit} = \lambda_{pi} + \lambda_{pt}$$

where $\lambda_{pt} \sim N(\lambda_p, w_p)$ and $\lambda_{pi} \sim N(0, w_i)$

- The random loadings accommodate measurement non-invariance across time and individual
- Model 4: Adding factor variance non-invariance across time. Can be done either by adding (a) introducing a factor model for the random loadings or (b) introducing a random loadings for the residual of the factor.
- We choose (b). $Var(f) = 0.51 + (0.7 + \sigma_t)^2$ where σ_t is a mean zero random effect

Aggressive-Disruptive Behavior Example Continued: Model 4 Setup

```

model:
%within%
s1-s9 | f by y1-y9;
s | f on time; f@0.51;
ss | e by f; e@1;

%between T1%
y1-y9;
s@0; [s@0];
s1-s9*1; [s1-s9*1];
ss*0.6; [ss@0.7];

%between ID%
y1-y9;
f*1; [f@0];
s*1; [s*0];
s1-s9*1; [s1-s9@0];
ss@0; [ss@0];

```

Aggressive-Disruptive Behavior Example Continued: Results For Categorical Analysis

Between ID Level					
Variances					
Y1	0.113	0.039	0.000	0.045	0.195
Y9	0.194	0.049	0.000	0.108	0.299
S1	0.082	0.029	0.000	0.041	0.152
S9	0.030	0.025	0.000	0.001	0.089
F	1.789	0.241	0.000	1.353	2.300
Between T1 Level					
Means					
S1	0.821	0.094	0.000	0.647	1.018
S9	1.049	0.125	0.000	0.809	1.299
Thresholds					
Y1\$1	-0.864	0.156	0.000	-1.183	-0.558
Y2\$1	-0.800	0.120	0.000	-1.041	-0.568
Variances					
Y1	0.120	0.231	0.000	0.035	0.666
Y9	0.073	0.158	0.000	0.019	0.416
S1	0.023	0.055	0.000	0.005	0.137
S9	0.050	0.110	0.000	0.011	0.302

Aggressive-Disruptive Behavior Example Conclusions

- Other extensions of the above model are possible, for example the growth trend can have time specific random effects: f and s can be free over time
- The more clusters there are on a particular level the more elaborate the model can be on that level. However, the more elaborate the model on a particular level is, the slower the convergence
- The main factor f can have a random effect on each of the levels, however the residuals Y_i should be uncorrelated on that level. If they are correlated through another factor model such as, fb by $y_1 - y_9$, then f would be confounded with that factor fb and the model will be poorly identified
- On each level the most general model would be (if there are no random slopes) the unconstrained variance covariance for the dependent variables Y_i . Any model that is a restriction of that model is in principle identified

Aggressive-Disruptive Behavior Example Conclusions, Continued

- Unlike ML and WLS multivariate modeling, for the time intensive Bayes cross-classified SEM, the more time points there are the more stable and easy to estimate the model is
- Bayesian methods solve problems not feasible with ML or WLS
- Time intensive data naturally fits in the cross-classified modeling framework
- Asparouhov and Muthén (2012). General Random Effect Latent Variable Modeling: Random Subjects, Items, Contexts, and Parameters

Cross-Classified / Multiple Membership Applications

- Jeon & Rabe-Hesketh (2012). Profile-Likelihood Approach for Estimating Generalized Linear Mixed Models With Factor Structures. JEBS
- Longitudinal growth model for student self-esteem
- Each student has 4 observations: 2 in middle school in wave 1 and 2, and 2 in high school in wave 3 and 4
- Students have multiple membership: Membership in middle school and in high school with a random effect from both
- Y_{ismh} is observation at time t for student s in middle school m and high school h

Cross-Classified / Multiple Membership Applications


- The model is

$$Y_{ismh} = \beta_1 + \beta_2 T2 + \beta_3 T3 + \beta_4 T4 + \delta_s + \delta_m \mu_t + \delta_h \lambda_t + \varepsilon_{ismh}$$

where T2, T3, T4 are dummy variables for wave 2, 3, 4

- δ_s , δ_m and δ_h are zero mean random effect contributions from student, middle school and high school
- $\mu_t = (1, \mu_2, \mu_3, \mu_4)$
- $\lambda_t = (0, 0, 1, \lambda_4)$, i.e., no contribution from the high school in wave 1 and 2 because the student is still in middle school
- ε_{ismh} is the residual
- Very simple to setup in Mplus

Cross-Classified / Multiple Membership Applications



Cross-Classified / Multiple Membership Applications

MODEL:

```
%WITHIN%
fs BY y1-y4@1;
[y1-y4];

%BETWEEN mschool%
fm BY y1@1 y2-y4;
y1-y4@0; [y1-y4@0];

%BETWEEN hschool%
fh BY y1@0 y2@0 y3@1 y4;
y1-y4@0; [y1-y4@0];
```